



## Study Manual for SOA Exam FAM-S 1<sup>st</sup> Edition, 2<sup>nd</sup> Printing

I.E.ARWTORN, I.E.AD TOMORROW

by Sam A. Broverman Ph.D., ASA

Your Integrated Study Manual Program Includes:





# Study Manual for SOA Exam FAM-S 1<sup>st</sup> Edition, 2<sup>nd</sup> Printing

by Sam A. Broverman Ph.D., ASA



Actuarial & Financial Risk Resource Materials Since 1972

Copyright © 2022, ACTEX Learning, a division of ArchiMedia Advantage Inc.

Printed in the United States of America.

No portion of this ACTEX Study Manual may be reproduced or transmitted in any part or by any means without the permission of the publisher.



Actuarial University is a reimagined platform built around a more simplified way to study. It combines all the products you use to study into one interactive learning center.



You can find integrated topics using this network icon. When this icon appears, it will be next to an important topic in the manual! Click the link in your digital manual, or search the underlined topic in your print manual.

## 1. Login to: www.actuarialuniversity.com

- 2. Locate the **Topic Search** on your exam dashboard, and enter the word or phrase into the search field, selecting the best match.
- 3. A topic "Hub" will display a list of integrated products that offer more ways to study the material.
- 4. Here is an example of the topic **Pareto Distribution**:

Pareto Distribution 
$$\checkmark$$
  
The (Type II) Pareto distribution with parameters  $\alpha, \beta > 0$  has pdf  
 $f(x) = \frac{\alpha\beta^{\alpha}}{(x+\beta)^{\alpha+1}}, \quad x > 0$   
and cdf  
 $F_P(x) = 1 - \left(\frac{\beta}{x+\beta}\right)^{\alpha}, \quad x > 0.$   
If X is Type II Pareto with parameters  $\alpha, \beta$  then  
 $E[X] = \frac{\beta}{\alpha-1}$  if  $\alpha > 1$ ,  
and  
 $V\alpha r[X] = \frac{\alpha\beta^2}{\alpha-2} - \left(\frac{\alpha\beta}{\alpha-1}\right)^2$  if  $\alpha > 2.$   
COAL for SRM  
ACTEX Manual for P  
Probability for Risk Management, 3rd Edition  
ASM Manual for IFM  
Related Topics  $\sim$ 

Within the **Hub**, there will be unlocked and locked products.

Unlocked Products are the products that you own.



Many of Actuarial University's features are already unlocked with your study program, including:

GOAL Practice Tool Virtual Flashcards Instructional Videos\*

Formula & Review Sheet

## **GOAL: Guided Online Actuarial Learning**

#### Your Adaptable, Customizable, Online Actuarial Exam Prep Tool.

GOAL is an eLearning test preparation tool for students to practice skills learned in class or from independent study. The online platform offers a massive database of SOA & CAS exam-style problems with detailed solutions.

**GOAL** offers **instructor support** as well as **performance reporting and tracking** to monitor your progress. Don't forget to check out **GOAL Score** to see how ready you are to sit for your exam!

#### Practice. Quiz. Test. Pass!

- 10,000+ Exam-style problems with detailed solutions!
- Adaptive Quizzes
- 3 Learning Modes
- 3 Difficulty Modes



\*Available standalone, or included with the Study Manual Program Video Bundle



## Free with your ACTEX or ASM Study Manual Program.

Available now for P, FM, FAM, ALTAM, ASTAM, SRM, MAS-I, MAS-II, CAS 6C/U

Prepare for your exam confidently with **GOAL** custom **Practice Sessions**, **Quizzes**, and **Simulated Exams**.

Actuarial University         QUESTION 14 OF 62       Question #	Prev Next		Quickly access the Hub for additional learning
Question	Difficulty: Mastery 🕄	,	
At time $t = 0$ year, Donald puts \$1,000 into a fund crediting is compounded semiannually. At time $t = 2$ years, Lewis puts \$1,000 into a different fund cr $\delta_t = 1/(5+t)$ for all $t$ .	nterest at a nominal rate of $i$ editing interest at a force		Flag problems for review, record notes, and email your professor.
At time $t = 10$ years, the amounts in each fund will be equal.			
Calculate <i>i</i> . Possible Answers			View difficulty level.
<ul> <li>★ 6.9%</li> <li>▼ 7.0%</li> <li>C 7.1%</li> <li>Help Me Start</li> </ul>	B 7.2% A 7.3%		Helpful strategies to get you
Equate the expressions for the AVs at $t = 16$ . Then solve for $i$	(2):	— (	started.
Solution Equate the expressions for for the AVs at $t = 16$ and calculate $(1 + i^{(2)}/2)^{32} = 3$ $(1 + i^{(2)}/2) = 3^{(1/32)} = 1.03493$	ĵ(2): ► 0:08/0:59		Full solutions, with detailed explanations to deepen your
$i^{(2)}/2 = 0.03493$	Donald: $a(16) = (1+i^{(2)}/2)^{2\cdot 16} = (1+i^{(2)}/2)^{32}$	l	understanding.
$i^{(2)}=7.0\%$	Lewis: $a(16) = e^{\int_{2}^{16} \frac{1}{6\pi t} dt} = e^{\ln(5+t)\int_{2}^{16}} = e^{\ln(21) - \ln(7)} = 21/7 = 3$		
Common Questions & Errors <u>Student Question 1:</u> After solving this problem I got .069855. A <u>Answer:</u> The provided answer choices are all rounded to 1 deci	Are we expected to round to .07? mal place. So the answer 6.9855%		Commonly encountered errors.
should be rounded to 7.0% to be correct to 1 decimal place. Rate this problem	nprovement 🦃 Inadequate		Rate a problem or give feedback



# Track your exam readiness with GOAL Score!

Available now for P, FM, FAM, ALTAM, ASTAM, SRM, MAS-I, MAS-II, CAS 6C/U

**GOAL Score** tracks your performance through GOAL Practice Sessions, Quizzes, and Exams, resulting in an aggregate, weighted score that gauges your exam preparedness.

By measuring both **your performance**, and **the consistency of your performance**, **GOAL Score** produces a reliable number that will give you confidence in your preparation before you sit for your exam.



## Contents

#### INTRODUCTORY COMMENTS

Section 1	Review of Probability	1
1.1	Basic Probability Concepts	1
1.2	Conditional Probability and Independence of Events	4
Section 2	Review of Random Variables I	19
2.1	Calculus Review	19
2.2	Discrete Random Variable	20
2.3	Continuous Random Variable	21
2.4	Mixed Distribution	23
2.5	Cumulative Distribution, Survival and Hazard Functions	23
2.6	Examples of Distribution Functions	24
2.7	The Empirical Distribution	27
2.8	Gamma Function and Related Functions	28
Section 3	Review of Random Variables II	35
3.1	Expected Value and Other Moments of a Random Variable	35
3.2	Percentiles and Quantiles of a Distribution	38
3.3	The Mode of a Distribution	38
3.4	Random Samples and The Sampling Distribution	40
3.5	The Normal Distribution	41
3.6	Approximating a Distribution Using a Normal Distribution	43
3.7	Distribution of a Transformation of Random Variable $X$	45
Section 4	Review of Random Variables III	<b>53</b>
4.1	Joint Distribution of Random Variables $X$ and $Y$	53
4.2	Marginal distribution of X found from a joint distribution of X and Y $\ldots$	56
4.3	Independence of Random Variables $X$ and $Y$	58
4.4	Conditional Distribution of Y Given $X = x$	58
4.5	Covariance Between Random Variables $X$ and $Y$	61
4.6	Coefficient of correlation between random variables $X$ and $Y$	62
Section 5	Parametric Distributions	69
5.1	Parametric Distributions	69
5.2	Families of Distributions	71

#### CONTENTS

5.2.1	1 The Linear Exponential Family 71
5.3	The Single Parameter Pareto and Two Parameter Pareto Distributions 72
5.4	A Note on the Poisson Distribution
Section 6	Distribution Tail Behavior 81
6.1	Measuring Tail Weight Using Existence of Moments
6.2	Comparing the Tail Weights of Two Distributions
Section 7	Mixture Of Two Distributions 85
7.1	Mixture of Two Distributions
7.2	Formulating a Mixture Distribution as a Combination of
	Conditional Distributions
7.3	The Variance of a Mixed Distribution IS (usually) NOT
	the Weighted Average of the Variances
Section 8	Mixture Of <i>n</i> Distributions 101
8.1	Mixture of $n$ Distributions $\ldots \ldots \ldots$
8.2	Two Important Rules of Probability
Section 9	Frequency Models The Number Of Claims 113
9.1	Poisson Distribution
9.2	Binomial Distribution
9.3	Negative Binomial Distribution
9.4	The $(a,b,0)$ Class of Discrete Distributions $\ldots \ldots \ldots$
9.5	The $(a,b,1)$ Class of Discrete Distributions $\ldots \ldots \ldots$
Section 10	Policy Limits And The Limited Loss 139
10.1	Policy Limit $u$ and the Limited Loss Random Variable $\ldots \ldots \ldots$
10.2	Limited Expected Value
Section 11	Policy Deductible I - The Cost Per Loss 147
11.1	Ordinary Policy Deductible $d$ and Cost Per Loss $\ldots \ldots \ldots \ldots \ldots \ldots 147$
11.2	Modeling Bonus Payments
11.3	Franchise Deductible
Section 12	Policy Deductible II - The Cost Per Payment 167
12.1	Ordinary Policy Deductible $d$ and Cost Per Payment $\ldots \ldots \ldots$
12.2	Expected Cost Per Payment
12.3	Variance of Cost Per Payment With Deductible $d$
12.4	Franchise Deductible $d$

Section 13	Deductibles Applied To The Uniform, Exponential And Pareto Distributions	189
Section 14	Combined Limit And Deductible	203
14.1	Maximum Covered Loss $u$ Combined With Policy Deductible $d < u$	. 203
14.2	PDF and CDF of $Y_L$ and $Y_P$	. 207
14.3	A Few Comments	. 208
14.4	Graphical Representation of $E[Y_L]$	. 209
Section 15	Additional Policy Adjustments	227
15.1	Coinsurance and Copay	. 227
15.2	Inflation	. 229
15.3	Policy Deductible $d$ in Combination With Maximum Covered Loss $u$ and Coinsurance Factor $\alpha$ and Inflation Factor $r$	. 231
Section 16	Aggregate Models - Compound Distributions	<b>243</b>
16.1	The Compound Distribution Model for Aggregate Claims in One Period of	049
16.9	The Course and Deissen Distribution	. 243
10.2	The Normal Approximation to a Compound Distribution	· 240
16.4	The Convolution Method for Finding the Distribution	. 240
	of a Sum of Random Variables	. 247
Section 17	More Properties Of The Aggregate Loss	<b>2</b> 81
17.1	The Individual Risk Model	. 281
Section 18	Stop Loss Insurance	299
Section 19	Risk Measures	319
19.1	Value at Risk	. 319
19.2	Tail-Value-at-Risk	. 320
Section 20	Parametric Models Data And Estimation Review	329
20.1	Review of Parametric Distributions	. 329
20.2	Data and Estimation	. 330
Section 21	Maximum Likelihood Estimation Based On Complete Data	333
Section 22	Maximum Likelihood Estimation Based On Incomplete Data	347
22.1	MLE Based on Interval Grouped Data	. 347
22.2	Maximum Likelihood Estimation Based On Censored and/or Truncated Dat	a349;
Section 23	MLE For The Exponential Distribution	365

#### CONTENTS

23.1	MLE of the Exponential Distribution Based on Complete Data
23.2	MLE of the Exponential Distribution
	Based on Policy Limit (Right-Censored) Data
23.3	MLE of the Exponential Distribution Based on Policy Deductible Data 366
23.4	MLE of the Exponential Distribution
	Based on General Policy Limit and Deductible Data
23.5	Interval Grouped Data and MLE Estimation of the Exponential Distribution 369
Section 24	MLE For Pareto And Weibull Distributions 383
24.1	Maximum Likelihood Estimation and Transformations
24.2	Pareto Distribution $\alpha$ , $\theta$ , where $\theta$ is given $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots 385$
24.3	Single Parameter Pareto Distribution, $\alpha, \theta$ , where $\theta$ is given $\ldots \ldots \ldots 389$
24.4	Weibull Distribution, $\tau, \theta$ , where $\tau$ is given
Section 25	MLE Applied To FAM Table Distributions 407
25.1	Inverse Exponential Distribution, $\theta$
25.2	Inverse Pareto Distribution, $\tau$ , $\theta$ , where $\theta$ is given
25.3	Inverse Weibull Distribution, $\tau$ , $\theta$ , where $\tau$ is given $\ldots \ldots \ldots \ldots \ldots \ldots 410$
25.4	Normal Distribution, $\mu$ , $\sigma^2$
25.5	Lognormal Distribution, $\mu$ , $\sigma^2$
25.6	Gamma Distribution, $\alpha$ , $\theta$ , where $\alpha$ is given $\ldots \ldots 413$
25.7	Inverse Gamma Distribution, $\alpha$ , $\theta$ , where $\alpha$ is given
25.8	poisson Distribution $\lambda$
25.9	Binomial Distribution $m, q$
25.10	Negative Binomial Distribution
Section 26	Limited Fluctuation Credibility 429
26.1	Introductory Comments on Credibility Theory
26.2	The Standard for Full Credibility
26.3	Full credibility applied to a frequency distribution
26.4	The Standard for Full Credibility Applied to Compound Distributions $\dots$ 436
26.5	Standard for Full Credibility Applied to Poisson Random Variable $N$ 438
26.6	Standard For Full Credibility Applied to a Compound Poisson Distribution 438
26.7	Partial Credibility
Section 27	Insurance And Reinsurance Coverages, Major Medical And Den-
	tal Coverage 467
27.1	Major Medical Coverages
27.2	Dental Coverage
Section 28	Pricing and Reserving Short-Term Insurance 471

#### CONTENTS

Section 29	Short-Term Insurance Loss Reserving 48	5
29.1	Basic Elements of Loss Reserving	35
29.2	Case Reserves Plus and Expected Loss Ratio Methods	35
29.3	Chain-Ladder (Loss Development Triangle) Method	36
29.4	Bornhuetter Ferguson Method	91
29.5	Discounted Reserves	92
29.6	Additional Notes and Comments On Chapter 3 of the Text 49	)3
Section 30	Short-Term Insurance Ratemaking 51	.5
30.1	Objectives of Ratemaking	15
30.2	Data for Ratemaking	15
30.3	Loss-Development and Trend Factors	16
30.4	Expenses, Profit and Contingency Loading	18
30.5	Credibility Factors	9
30.6	Overall Rate Changes	9
30.7	Rate Classification Differentials	21
Section 31	Reinsurance 52	29
Section 32	Option Pricing 53	35
32.1	Arbitrage	35
32.2	Introduction to Options	36
32.3	The Binomial Option Pricing Model For One Period	38
32.4	The Binomial Option Pricing Model For Two Periods	13
32.5	The Black-Scholes-Merton Option Pricing Model	19
Practice Ex	cam 1 55	59
Solutions to	Practice Exam 1	36
Practice Ex	xam 2 57	'3
Solutions to	Practice Exam 2	79
Practice Ex	xam 3 58	37
Solutions to	Practice Exam 3	<b>)</b> 4
Practice Ex	cam 4 60	)3
Solutions to	Practice Exam 4	10
Index	61	.9

## **INTRODUCTORY COMMENTS**

The FAM exam is divided almost equally into FAM-S and FAM-L topics. This study guide is designed to help in the preparation for the Society of Actuaries FAM-S Exam.

The first part of this manual consists of a summary of notes, illustrative examples and problem sets with detailed solutions. The second part consists of 4 practice exams. The SOA exam syllabus for the FAM exam indicates that the exam is 3.5 hours in length with 40 multiple choice questions. The practice exams in this manual each have 20 questions, reflecting the fact that FAM-S is 50% of the full FAM exam. The appropriate time for the 20 question FAM-S practice exams in this manual is one hour and forty-five minutes.

The level of difficulty of the practice exam questions has been designed to be similar to those on past exams covering the same topics. The practice exam questions are not from old SOA exams.

I have attempted to be thorough in the coverage of the topics upon which the exam is based, and consistent with the notation and content of the official references. I have been, perhaps, more thorough than necessary on reviewing maximum likelihood estimation.

Because of the time constraint on the exam, a crucial aspect of exam taking is the ability to work quickly. I believe that working through many problems and examples is a good way to build up the speed at which you work. It can also be worthwhile to work through problems that you have been done before, as this helps to reinforce familiarity, understanding and confidence. Working many problems will also help in being able to more quickly identify topic and question types. I have attempted, wherever possible, to emphasize shortcuts and efficient and systematic ways of setting up solutions. There are also occasional comments on interpretation of the language used in some exam questions. While the focus of the study guide is on exam preparation, from time to time there will be comments on underlying theory in places that I feel those comments may provide useful insight into a topic.

The notes and examples are divided into 32 sections of varying lengths, with some suggested time frames for covering the material. There are almost 180 examples in the notes and over 440 exercises in the problem sets, all with detailed solutions. The 4 practice exams have 20 questions each, also with detailed solutions. Some of the examples and exercises are taken from previous SOA exams. Some of the problem set exercises are more in depth than actual exam questions, but the practice exam questions have been created in an attempt to replicate the level of depth and difficulty of actual exam questions. In total there are almost 700 examples/problems/sample exam questions with detailed solutions. ACTEX gratefully acknowledges the SOA for allowing the use of their exam problems in this study guide.

I suggest that you work through the study guide by studying a section of notes and then attempting the exercises in the problem set that follows that section. The order of the sections of notes is the order that I recommend in covering the material, although the material on pricing and reserving in Sections 27 to 31 is independent of the other material on the exam. The order of topics in this manual is not the same as the order presented on the exam syllabus.

It has been my intention to make this study guide self-contained and comprehensive for the FAM-S Exam topics, however it is important to be familiar with original reference material on all topics.

While the ability to derive formulas used on the exam is usually not the focus of an exam question, it is useful in enhancing the understanding of the material and may be helpful in memorizing formulas. There may be an occasional reference in the review notes to a derivation, but you are encouraged to review the official reference material for more detail on formula derivations.

In order for the review notes in this study guide to be most effective, you should have some background at the junior or senior college level in probability and statistics. It will be assumed that you are reasonably familiar with differential and integral calculus. The prerequisite concepts to modeling and model estimation are reviewed in this study guide. The study guide begins with a detailed review of probability distribution concepts such as distribution function, hazard rate, expectation and variance. Of the various calculators that are allowed for use on the exam, I am most familiar with the BA II PLUS. It has several easily accessible memories. The TI-30X IIS has the advantage of a multi-line display. Both have the functionality needed for the exam.

There is a set of tables that has been provided with the exam in past sittings. These tables consist of some detailed description of a number of probability distributions along with tables for the standard normal and chi-squared distributions. The tables can be downloaded from the SOA website www.soa.org.

If you have any questions, comments, criticisms or compliments regarding this study guide, please contact the publisher ACTEX, or you may contact me directly at the address below. I apologize in advance for any errors, typographical or otherwise, that you might find, and it would be greatly appreciated if you would bring them to my attention. ACTEX will be maintaining a website for errata that can be accessed from www.actexmadriver.com. It is my sincere hope that you find this study guide helpful and useful in your preparation for the exam. I wish you the best of luck on the exam.

Samuel A. Broverman			July 2022
Department of Statistical	Sciences	www	.sambroverman.com
University of Toronto	E-mail: sam.broverman@utoronto.ca	or	2brove@rogers.com

### Section 7

## Mixture Of Two Distributions

The material in this section relates to Section 4.2.3 of "Loss Models". The suggested time frame for this section is 2 hours. The topic of distribution mixtures is not mentioned in the learning objectives for the FAM exam but all of Chapter 4 of "Loss Models" is listed in the reference reading.

#### 7.1 Mixture of Two Distributions

We begin with a formal algebraic definition of how a mixture of two distributions is constructed, and later we will look at how a mixture distribution is described by general reasoning.

Given random variables  $X_1$  and  $X_2$ , with pdf's or pf's  $f_{X_1}(x)$  and  $f_{X_2}(x)$ , and given 0 < a < 1, we construct the random variable Y with pdf

$$f_Y(y) = a \times f_{X_1}(y) + (1-a) \times f_{X_2}(y)$$
(7.1)

Y is called a mixture distribution or a two-point mixture of the distributions of  $X_1$  and  $X_2$ .

The two-point mixture random variable Y can also be defined in terms of the cdf,

$$F_{Y}(y) = a \times F_{X_{1}}(y) + (1 - a) \times F_{X_{2}}(y)$$
(7.2)

 $X_1$  and  $X_2$  are the **component distributions** of the mixture, and the factors a and 1-a are referred to as **mixing weights**. It is important to understand that we are **not adding**  $aX_1$  and  $(1-a)X_2$ , Y **is not** a sum of random variables. Y is defined in terms of a pdf (or cdf) that is a weighted average of the pdf's (or cdf's) of  $X_1$  and  $X_2$ . We are adding  $af_{X_1}$  and  $(1-a)f_{X_2}$  to get  $f_Y$ .

**Example 7.1.** As a simple illustration of a mixture distribution, consider two bowls. Bowl A has 5 balls with the number 1 on them and 5 balls with the number 2 on them, and bowl B has 3 balls with the number 1 and 7 balls with the number 2. Let  $X_1$  denote the number on a ball randomly chosen from bowl A, and let  $X_2$  denote the number on a ball randomly chosen from bowl B. The probability functions of  $X_1$  and  $X_2$  are  $f_{X_1}(1) = f_{X_1}(2) = 0.5$  and  $f_{X_2}(1) = 0.3$ ,  $f_{X_2}(2) = 0.7$ .

Suppose we create the mixture distribution with mixing weights a = 0.5 and 1 - a = 0.5. The mixture distribution Y has probability function  $f_Y(1) = 0.5 \times 0.5 + 0.5 \times 0.3 = 0.4$ ,  $f_Y(2) = 0.5 \times 0.5 + 0.5 \times 0.7 = 0.6$ .

Note that the outcomes (ball numbers) of the mixture distribution Y come from the possible outcomes of the component distributions  $X_1$  and  $X_2$ .

An alternative way of looking at this mixture distribution is by means of conditioning on a "parameter". This will be important when we look at continuous mixing. The parameter approach to describe the mixture distribution in Example 7.1 is as follows.

Suppose that a fair coin is tossed. If the toss is a head, a ball is chosen at random from bowl A and if the toss is a tail, a ball is chosen at random from bowl B. We define the random variable Z to be the number on the ball. We will see that Z has the same distribution as the mixture distribution labeled Y above.

The random variable Z can be interpreted as follows. Consider the 2-point random variable  $\Theta$ , for which  $\Theta =$ Bowl A if the coin toss is a head, and  $\Theta =$ Bowl B if the toss is a tail. Then  $P(\Theta = A) = P(\Theta = B) = .5$  (since the coin is fair).  $\Theta$  is used to indicate which bowl the ball will be chosen from depending on the outcome of the coin toss.

If the toss is a head, the bowl is A, and then Z has the  $X_1$  distribution for the number on the ball, so  $f_{X_1}(1) = P(Z = 1|\Theta = A) = 0.5$  and  $f_{X_1}(2) = P(Z = 2|\Theta = A) = 0.5$ . In a similar way, if the toss is a tail, the bowl is B, and then Z has the  $X_2$  distribution for the number on the ball, so  $f_{X_2}(1) = P(Z = 1|\Theta = B) = 0.3$  and  $f_{X_2}(2) = P(Z = 2|\Theta = B) = 0.7$ .

Z is described as a combination of two conditional distributions based on the parameter  $\Theta$ .

To find the overall, or unconditional distribution of Z, we use some basic rules of probability. Since  $\Theta$  must be A or B, we can think of bowl B as the "complement" of bowl A, and then

$$P(Z = 1) = P[(Z = 1) \cap (\Theta = A)] + P[(Z = 1) \cap (\Theta = B)]$$
  
=  $P(Z = 1|\Theta = A) \times P(\Theta = A) + P(Z = 1|\Theta = B) \times P(\Theta = B)$   
=  $0.5 \times 0.5 + 0.5 \times 0.3 = 0.4$ 

We have used the rule  $P(C) = P[C \cap D] + P[C \cap D'] = P(C|D) \times P(D) + P(C|D') \times P(D')$ .

This shows that the distribution of Z is the same as the mixture distribution Y in Example 7.1. The mixing weights for the two bowls are the probabilities of the coin indicating bowl A or bowl B.

#### Language used on exam questions that identifies a mixture distribution

There is some typical language that is used in exam questions that indicates that a mixture of distributions is being considered. This language will be illustrated using the distributions involved in the bowl example.

Suppose that we are told that there are two types of individuals. Type A individuals have a mortality probability of .5 (and survival probability of .5) in the coming year, and Type B individuals have a mortality probability of .3 in the coming year. In a large group of these individuals, 50% are Type A and 50% are Type B. An individual is chosen at random from the group. We want to find this randomly chosen individual's mortality probability.

We can use the usual rules of conditional probability to formulate this probability, just as above:

$$\begin{aligned} P(\text{dying this year}) &= P(\text{dying} \cap \text{Type A}) + P(\text{dying} \cap \text{Type B}) \\ &= P(\text{dying}|\text{Type A}) \times P(\text{Type A}) + P(\text{dying} \cap \text{Type B}) \times P(\text{Type B}) \\ &= 0.5 \times 0.5 + 0.5 \times 0.3 = 0.4. \end{aligned}$$

This is exactly  $f_Y(1) = f_{X_1}(1) \times a + f_{X_2}(1) \times (1-a)$  where "1" corresponds to the event of dying within the year. Note that the phrase "50% are Type A" is interpreted as meaning that if an individual is chosen from the large group, the probability of being Type A is 0.5. This is language that is often used in exam questions to indicate a mixture.

#### Mixture of a discrete and a continuous distribution

In Section 2 of this study guide we looked at "mixed distributions". In that section, a mixed distribution referred to a random variable that was continuous on part of its probability space and also had one or more discrete points. The concept of mixed distribution just introduced in this section can be used to describe the Section 2 type of mixed distribution. The following example uses the example from Section 2 to illustrate this.

**Example 7.2.** Suppose that X is defined in the following piecewise way: X has a discrete point of probability of .5 at X = 0, and on the interval (0,1) X is a continuous random variable with density function f(x) = x for 0 < x < 1, and X has no density or probability elsewhere. Show that this random variable can be described as a mixture of two distributions with appropriate definitions for component distributions  $X_1$ ,  $X_2$  and mixing weights a and 1 - a.

#### Solution.

Let  $X_1 = 0$  be a constant (not actually a random variable, but is sometimes called a degenerate random variable), so that  $f_{X_1}(0) = P(X_1 = 0) = 1$  and  $f_{X_1}(x) = 0$  for  $x \neq 0$ .

Let  $X_2$  be continuous on (0,1) with pdf  $f_{X_2}(x) = 2x$ .

With mixing weights of a = 0.5 and 1 - a = 0.5, using the definition of the mixture of two distributions, we have the mixed random variable Y satisfying

 $f_Y(0) = a \times f_{X_1}(0) + (1-a) \times f_{X_2}(0) = (0.5 \times 1) + 0 = 0.5, \text{ and}$   $f_Y(x) = a \times f_{X_1}(x) + (1-a) \times f_{X_2}(x) = 0.5 \times 0 + 0.5 \times 2x = x \text{ for } 0 < x < 1.$ *Y* has the same distribution as the original *X*.

We should be a little careful about the situation in Example 7.2. When we are mixing discrete probability points with a continuous density, for any particular discrete point we always assign a probability of 0 at that point for any continuous component distribution. For instance, in Example 7.2, if  $X_1$  was at the single point 0.4, then  $f_{X_1}(0.4) = 1$  and  $f_{X_1}(x) = 0$  elsewhere, and for the mixture distribution, then  $f_Y(0.4) = 0.5 \times 1 = 0.5$  (we do not add  $0.5 \times f_{X_2}(0.4)$ ).

Some important relationships for mixture distributions

We have already seen the defining relationships

$$f_Y(y) = a f_{X_1}(y) + (1-a) f_{X_2}(y)$$

and

$$F_Y(y) = aF_{X_1}(y) + (1-a)F_{X_2}(y).$$

We can interpret these relationships by saying that the pdf and cdf of Y are weighted averages of the component pdf's and cdf's. This weighted average interpretation can be applied to a number of other distribution related quantities.

- if C is any event related to Y, then  $P(C) = a \times P_{X_1}(C) + (1-a) \times P_{X_2}(C);$  (7.3)  $P_{X_1}(C)$  is the event probability based on random variable  $X_1$ , and the same for  $X_2$
- If g is any function (that doesn't involve the parameters of Y), then  $E[q(Y)] = a \times E[q(X_1)] + (1-a) \times E[q(X_2)]$ (7.4)

The justification for this relationship follows from the form of the mixed density;

$$\begin{split} E[g(Y)] &= \int g(y) \, f_Y(y) \, dy = \int g(y) \times [a \times f_{X_1}(y) + (1-a) \times f_{X_2}(y)] \, dy \\ &= a \times \int g(y) \times f_{X_1}(y) \, dy + (1-a) \times \int g(y) \times f_{X_2}(y) \, dy \\ &= a \times E[g(X_1)] + (1-a) \times E[g(X_2)] \end{split}$$

Some of the important examples of these weighted average relationships are:

Interval Probability: the event C is  $c < Y \leq d$ 

$$P(c < Y \le d) = a \times P(c < X_1 \le d) + (1 - a) \times P(c < X_2 \le d)$$
(7.5)

kth moment of Y: the function g is  $g(Y) = Y^k$ 

$$E[Y^{k}] = a \times E[X_{1}^{k}] + (1 - a) \times E[X_{2}^{k}]$$
(7.6)

#### 7.2 Formulating a Mixture Distribution as a Combination of Conditional Distributions

The relationships above can also be formulated as applications of probability rules involving conditional probability and conditional expectation.

(i) For any random variable Y and event D,

$$f_Y(y) = f(y|D) \times P(D) + f(y|D') \times P(D')$$
(7.7)

(ii) For any events C and D,

$$P[C] = P[C \cap D] + P[C \cap D'] = P[C|D] \times P[D] + P[C|D'] \times P[D'].$$
(7.8)

(iii) For any random variable Y and any event D,

$$E[Y] = E[Y|D] \times P[D] + E[Y|D'] \times P[D']$$
(7.9)

We can define the random variable  $\Theta$  to be 1 or 2, indicating whether or not event D has occurred, so  $\Theta = 1 \equiv D$  has occurred, and  $\Theta = 2 \equiv D'$  has not occurred.

We can think of  $X_1$  as the conditional distribution of Y given  $\Theta = 1$  (event D) and we can think of  $X_2$  as the conditional distribution of Y given  $\Theta = 2$  (event D'), so that

$$f(y|D) = f(y|\Theta = 1) = f_{X_1}(y)$$
 and  $f(y|D') = f(y|\Theta = 2) = f_{X_2}(y)$ .

The cdf, pdf and expectation of Y are adaptations of the conditional probability rules above. For instance, the mean of the mixed distribution is the "mixture of the means" of the component distributions.

$$E[Y] = E[Y|\Theta = 1] \times P[\Theta = 1] + E[Y|\Theta = 2] \times P[\Theta = 2] = E[X_1] \times a + E[X_2] \times (1 - a).$$

~

**Example 7.3.** A collection of insurance policies consists of two types. 25% of policies are Type 1 and 75% of policies are Type 2. For a policy of Type 1, the loss amount per year follows an exponential distribution with mean 200, and for a policy of Type 2, the loss amount per year follows a Pareto distribution with parameters  $\alpha = 3$  and  $\theta = 200$ . For a policy chosen at random from the entire collection of both types of policies, find the probability that the annual loss will be less than 100, and find the average loss.

#### Solution.

The two types of losses are the random variables  $X_1$  and  $X_2$ .

 $X_1$  has an exponential distribution with mean 200, so it has cdf  $F_{X_1}(x) = 1 - e^{-x/200}$ .

$$X_2$$
 has cdf  $F_{X_2}(x) = 1 - \left(\frac{\theta}{x+\theta}\right)^{\alpha} = 1 - \left(\frac{200}{x+200}\right)^3$ .

These are the component distributions. The mixing weights are the proportions of policies of each type, so that a = 0.25 (the proportion of Type 1 policies) and 1 - a = 0.75. The loss random variable Y of the randomly chosen policy has a distribution that is the mixture of  $X_1$  and  $X_2$ . In this context, "randomly chosen" is taken to mean that the probabilities of choosing a Type 1 or Type 2 policy are the proportions of those policy types in the full collection of policies.

The cdf of Y is

$$F_Y(y) = a \times F_{X_1}(y) + (1-a) \times F_{X_2}(y)$$
  
= 0.25 × (1 - e<sup>-y/200</sup>) + 0.75 ×  $\left(1 - \left(\frac{200}{y+200}\right)^3\right)$ 

Then,  $F_Y(100) = 0.626$ .

The mean of Y is  $0.25 \times E[X_1] + 0.75 \times E[X_2] = 0.25 \times 200 + 0.75 \times \frac{200}{3-1} = 125$ .

#### 

#### 7.3 The Variance of a Mixed Distribution IS (usually) NOT the Weighted Average of the Variances

For a mixture of two random variables, the weighted average pattern applies to most quantities for the mixed distribution.

One important exception to this rule is the formulation of the variance of the mixture. If  $Y \cdot P$  is any random variable, then we can formulate the variance of Y as  $Var[Y] = E[Y^2] - (E[Y])^2$ .

If Y is the mixture of  $X_1$  and  $X_2$  with mixing weights a and 1-a, then we know that  $E[Y] = a \times E[X_1] + (1-a) \times E[X_2]$  and  $E[Y^2] = a \times E[X_1^2] + (1-a) \times E[X_2^2]$ .

It is tempting to guess that Var[Y] is the weighted average of  $Var[X_1]$  and  $Var[X_2]$ .

This is <u>not true</u> in general (we will see a special case later for which it is true).

**Example 7.4.** Find the variance of the loss on the policy chosen at random in Example 7.3, and compare it to the weighted average of the variances of the two component loss distributions.

#### Solution.

The mean of Y is  $0.25 \times E[X_1] + 0.75 \times E[X_2] = 0.25 \times 200 + 0.75 \times \frac{200}{3-1} = 125$ , and the second moment of Y is  $0.25 \times E[X_1^2] + 0.75 \times E[X_2^2] = 0.25 \times 2 \times 200^2 + 0.75 \times \frac{200^2 \times 2}{(3-1)(3-2)} = 50,000$ , so  $Var[Y] = E[Y^2] - (E[Y])^2 = 34,375$ .  $Var[X_1] = 40,000$  (variance of an exponential distribution is the square of the mean).  $Var[X_2] = \frac{200^2 \times 2}{(3-1)(3-2)} - 100^2 = 30,000$ . The weighted average of the two variances is  $0.25 \times 40,000 + 0.75 \times 30,000 = 32,500$ , which is not the variance of Y.

Weighted average does not apply to percentiles in a mixture distribution In Example 7.3, the 50th percentile of  $X_1$  is  $m_1$ , the solution of the equation  $F_{X_1}(m_1) = 1 - e^{-m_1/200} = 0.5$ , so that  $m_1 = 138.63$ .

The 50th percentile of  $X_2$  is  $m_2$ , the solution of  $F_{X_2}(m_2) = 1 - \left(\frac{200}{m_2 + 200}\right)^3 = 0.5$ , so that  $m_2 = 51.98$ .

The weighted average of the two 50th percentiles is  $0.25 \times 138.63 + 0.75 \times 51.98 = 73.64$ . The cdf of the mixed distribution Y is  $F_Y(y) = 0.25 \times F_{X_1}(y) + 0.75 \times F_{X_2}(y)$ , and  $F_Y(73.64) = 0.53$ , so 73.64 is not the 50th percentile of Y.

To find the 50th percentile, say m, of the mixed distribution in Example 7.3, we must solve the equation

$$50 = F_Y(m) = 0.25 \times F_{X_1}(m) + 0.75 \times F_{X_2}(m)$$
$$= 0.25 \times (1 - e^{-m/200}) + 0.75 \times \left(1 - \left(\frac{200}{m + 200}\right)^3\right).$$

There is no algebraic solution, and m would have to be found by a numerical approximation method. The 50th percentile turns out to be about 65.7.

#### Section 7 Problem Set

#### Mixture Of Two Distributions

- 1. A portfolio of insurance policies is divided into low and high risk policies. 80% of the policies are low risk and the rest are high risk. The annual number of claims on a low risk policy has a Poisson distribution with a mean of .25 and the annual number of claims on a high risk policy has a Poisson distribution with a mean of 2. A policy is randomly chosen from the portfolio.
  - (a) Find the mean and variance of the number of annual claims for the policy.
  - (b) Find the probability that the policy has at most 1 claim in the coming year.
- 2.  $\checkmark$  The distribution of a loss, X, is a two-point mixture:
  - (i) With probability 0.8, X has a two-parameter Pareto distribution with  $\alpha=2$  and  $\theta=100.$
  - (ii) With probability 0.2, X has a two-parameter Pareto distribution with  $\alpha=4$  and  $\theta=3000.$

Calculate  $Pr(X \le 200)$ .

$$(A) 0.76 (B) 0.79 (C) 0.82 (D) 0.85 (E) 0.88 (E) 0.88$$

3.  $\checkmark$  The random variable N has a mixed distribution:

- (i) With probability p, N has a binomial distribution with q = 0.5 and m = 2.
- (ii) With probability 1 p, N has a binomial distribution with q = 0.5 and m = 4.

Which of the following is a correct expression for  $\operatorname{Prob}(N=2)$ ?

- (A)  $0.125p^2$  (B) 0.375 + 0.125p
- (C)  $0.375 + 0.125p^2$  (D)  $0.375 0.125p^2$
- (E) 0.375-0.125p

4. • Y is a mixture of two exponential distributions,  $f_Y(y) = \frac{1}{2}e^{-y} + \frac{1}{6}e^{-y/3}$ . The random variable Z defined by the equation Z = 2Y. Z is a mixture of two exponentials. The means of those two exponential distributions are

- (A) 1 and 3 (B) 1 and 6
- (C) 2 and 3 (D) 2 and 6
- (E) 3 and 6
- 5.  $\checkmark$   $X_1$  has a uniform distribution on the interval (0,1000) and  $X_2$  has a uniform distribution on the interval (0,2000). Y is defined as a mixture of  $X_1$  and  $X_2$  with mixing weights of .5 for each mixture component. Find the pdf, cdf and median (50th percentile) of Y.

- 6. Solution of people aged 50 consists of twice as many non-smokers as smokers. Non-smokers at age 50 have a mortality probability of .1 and smokers at age 50 have a mortality probability of .2. Two 50-year old individuals are chosen at random from the population.
  - (a) Find the probability that at least one of them dies before age 51.
  - (b) Suppose that the mortality probabilities for smokers and non-smokers remain the same at age 51. Find the mortality probability of a randomly chosen survivor at age 51 in this population.
- 7.  $\checkmark$  Y is the mixture of an exponential random variable with mean 1 and mixing weight  $\frac{2}{3}$ , and an exponential distribution with mean 2 and mixing weight  $\frac{1}{3}$ .

Find the pdf, cdf, mean, variance and 90th percentile of Y.

- 8. V Which of the following statements are true?
  - I. A mixture of two different exponentials with the mixture having a mean 2 has a heavier right tail than a single exponential distribution with mean 2.
  - II. If  $f_Y(y) = a_1 \times f_{X_1}(y) + a_2 \times f_{X_2}(y)$ , where  $0 < a_1 < 1$  and  $0 < a_2 < 1$ , then  $e_Y(d) = a_1 \times e_{X_1}(d) + a_2 \times e_{X_2}(d)$ .
- 9.  $\checkmark$  You are given two independent estimates of an unknown quantity,  $\mu$ :
  - (i) Estimate A:  $E(\mu_A) = 1000$  and  $\sigma(\mu_A) = 400$
  - (ii) Estimate B:  $E(\mu_B) = 1200$  and  $\sigma(\mu_B) = 200$

Estimate C is a weighted average of the two estimates A and B, such that:

$$\mu_C = w \times \mu_A + (1 - w) \times \mu_B$$

Determine the value of w that minimizes  $\sigma(\mu_C)$ .

- (A) 0 (B) 1/5 (C) 1/4 (D) 1/3 (E) 1/2
- 10. You are given the claim count data for which the sample mean is roughly equal to the sample variance. Thus you would like to use a claim count model that has its mean equal to its variance. An obvious choice is the Poisson distribution. Determine which of the following models may also be appropriate.
  - (A) A mixture of two binomial distributions with different means.
  - (B) A mixture of two Poisson distributions with different means.
  - (C) A mixture of two negative binomial distributions with different means.
  - (D) None of A, B, or C
  - (E) All of A, B, and C

- 11. Cosses come from an equally weighted mixture of an exponential distribution with mean  $m_1$ , and an exponential distribution with mean  $m_2$ . Determine the least upper bound for the coefficient of variation of this distribution.
  - (A) 1 (B)  $\sqrt{2}$  (C)  $\sqrt{3}$  (D) 2 (E)  $\sqrt{5}$

12. X has the following distribution : Pr[X = 0] = 0.4, Pr[X = 1] = 0.6. The distribution of Y is conditional on the value of X: if X = 0, then the distribution of Y is Pr[Y = 0] = 0.6, Pr[Y = 1] = 0.2, Pr[Y = 2] = 0.2, and if X = 1, then the distribution of Y is Pr[Y = 0] = 0.2, Pr[Y = 1] = 0.3, Pr[Y = 2] = 0.5. Z is the sum of Y independent normal random variables, each with mean and variance 2.

What is Var[Z]?

- (A) 5.0 (B) 6.0 (C) 7.0 (D) 8.0 (E) 9.0
- 13. Subway trains arrive at your station at a Poisson rate of 20 per hour. 25% of the trains are express and 75% are local. The types and number of trains arriving are independent. An express gets you to work in 16 minutes and a local gets you there in 28 minutes. You always take the first train to arrive. Your co-worker always takes the first express. You are both waiting at the same station.

Calculate the conditional probability that you arrive at work before your co-worker, given that a local arrives first.

(A) 
$$37\%$$
 (B)  $40\%$  (C)  $43\%$  (D)  $46\%$  (E)  $49\%$ 

#### Section 7 Problem Set Solutions

1. (a) Y is the mixture of two Poisson distributions.  $E[Y] = 0.8 \times E[X_L] + 0.2 \times E[X_H] = 0.8 \times 0.250 + 0.2 \times 2 = 0.6$ . To find the second moment of Y, we recall that for a Poisson distribution with mean  $\lambda$ , the variance is also  $\lambda$ . Since  $Var[X] = E[X^2] - (E[X])^2$ , it follows that  $E[X^2] = Var[X] + (E[X])^2 = \lambda + \lambda^2$  for a Poisson distribution. Then,  $E[Y^2] = 0.8 \times E[X_L^2] + 0.2 \times E[X_H^2] = 0.8 \times (0.25 + 0.25^2) + 0.2 \times (2 + 2^2) = 1.45$ . Then  $Var[Y] = 1.45 - (0.6)^2 = 1.09$ .

(b) 
$$P(Y \le 1) = 0.8 \times P(X_1 \le 1) + 0.2 \times P(X_2 \le 1)$$
  
=  $0.8 \times [e^{-0.25} + 0.25e^{-0.25}] + 0.2 \times [e^{-2} + 2e^{-2}] = 0.860.$ 

2. The probability is a mixture of the probabilities for the two components.  $P(X \le 200) = 0.8 \times P(X_1 \le 200) + 0.2 \times P(X_2 \le 200)$ , where  $X_1$  and  $X_2$  are the two Pareto distributions.

$$P(X_1 \le 200) = 1 - \left(\frac{100}{200 + 100}\right)^2 = .8889, \text{ and}$$
  

$$P(X_2 \le 200) = 1 - \left(\frac{3000}{200 + 3000}\right)^4 = .2275.$$
  

$$P(X \le 200) = (0.8 \times 0.8889) + (0.2 \times 0.2275) = 0.757.$$

Answer A

3. 
$$P(N = 2) = p \times P(N_1 = 2) + (1 - p) \times P(N_2 = 2)$$
  
=  $p \times (0.5)^2 + (1 - p) \times 6 \times (0.5)^4 = 0.375 - 0.125p.$ 

We have used the binomial probabilities of the form  $\binom{m}{k}q^k(1-q)^{m-k}$ .

Answer E

4. 
$$f_Y(y) = \frac{1}{2} \times e^{-y} + \frac{1}{2} \times \frac{1}{3} e^{-y/3}$$
.  $F_Y(y) = \frac{1}{2} \times (1 - e^{-y}) + \frac{1}{2} \times (1 - e^{-y/3})$   
 $F_Z(z) = F_Y(\frac{z}{2}) = \frac{1}{2} \times (1 - e^{-z/2}) + \frac{1}{2} \times (1 - e^{-z/6})$ .  
Z is a mixture of two exponentials with means 2 and 6.

Answer D

5. 
$$f_Y(y) = 0.5 \times f_{X_1}(y) + 0.5 \times f_{X_2}(y)$$
  

$$= \begin{cases} 0.5 \times 0.001 + 0.5 \times 0.0005 = 0.00075 & \text{if } 0 < y < 1000 \\ 0.5 \times 0.0005 = 0.00025 & \text{if } 1000 \le y < 2000 \end{cases}$$
 $F_Y(y) = 0.5 \times F_{X_1}(y) + 0.5 \times F_{X_2}(y)$   

$$= \begin{cases} 0.5 \times 0.001x + 0.5 \times 0.0005x = 0.00075x & \text{if } 0 < y < 1000 \\ 0.5 \times 1 + 0.5 \times 0.0005x = 0.5 + 0.00025x & \text{if } 1000 \le y < 2000 \end{cases}$$

The median of Y, m, must satisfy the equation  $F_Y(m) = 0.5$ . We see that at y = 1000,  $F_Y(1000) = 0.75$ . Therefore, m < 1000, so that  $F_Y(m) = 0.00075m = 0.5$ . Therefore, m = 666.67. Note that the mean of Y is  $0.5 \times 500 + 0.5 \times 1000 = 750$ . 6. For a randomly chosen individual at age 50, the mortality probability is the mixture of the mortality probabilities for non-smokers and smokers. This is  $q = \frac{2}{3} \times 0.1 + \frac{1}{3} \times 0.2 = \frac{2}{15}$ .

The probability that both of two independent 50-year old individuals survive the year is  $(1 - \frac{2}{15})^2 = 0.7511$ . The probability at least one of them dies by age 51 is 1 - 0.7511 = 0.25.

Suppose that there are 1000 non-smokers and 500 smokers at age 50. The expected number of surviving non-smokers at age 51 is  $1000 \times 0.9 = 900$ , and the number of surviving smokers is  $500 \times 0.8 = 400$ .

A randomly chosen survivor at age 51 has a  $\frac{9}{13}$  chance of being a non-smoker and a  $\frac{4}{13}$  chance of being a smoker.

The mortality probability at age 51 for the randomly chosen survivor is  $\frac{9}{13} \times 0.1 + \frac{4}{13} \times 0.2 = .131$ .

7.  $f_Y(y) = \frac{2}{3} \times e^{-y} + \frac{1}{3} \times \frac{1}{2} e^{-y/2}$ .  $F_Y(y) = \frac{2}{3} \times (1 - e^{-y}) + \frac{1}{3} \times (1 - e^{-y/2})$ .  $E[Y] = \frac{2}{3} \times 1 + \frac{1}{3} \times 2 = \frac{4}{3}$ .  $E[Y^2] = \frac{2}{3} \times (2 \times 1^2) + \frac{1}{3} \times (2 \times 2^2) = 4 \rightarrow Var[Y] = 4 - (\frac{4}{3})^2 = \frac{20}{9}$ .

The 90th percentile of Y is c, which must satisfy the equation  $F_Y(c) = \frac{2}{3} \times (1 - e^{-c}) + \frac{1}{3} \times (1 - e^{-c/2}) = 0.9.$ 

If we let  $r = e^{-c/2}$ , then this becomes a quadratic equation in r,  $\frac{2}{3} \times (1 - r^2) + \frac{1}{3} \times (1 - r) = 0.9$ , or equivalently,  $2r^2 + r - .3 = 0$ .

Solving for r results in r = .21 or r = -0.71.

We ignore the negative root, since  $r = e^{-c/2}$  must be positive.

Then  $c = -2 \ln r = -2 \ln .21 = 3.12$  is the 90th percentile of Y.

8. I. If  $f_Y(y) = a_1 \times f_{X_1}(y) + a_2 \times f_{X_2}(y) = a_1 \times \frac{1}{\theta_1} e^{-y/\theta_1} + a_2 \times \frac{1}{\theta_2} e^{-y/\theta_2}$ , and  $a_1\theta_1 + a_2\theta_2 = 2$ . Then  $\theta_1 < 2 < \theta_2$  (or vice-versa).

For an exponential distribution with mean 2, we have  $f_Z(y) = \frac{1}{2}e^{-y/2}$   $\frac{f_Y(y)}{f_Z(y)} = \frac{a_1 \times \frac{1}{\theta_1} e^{-y/\theta_1} + a_2 \times \frac{1}{\theta_2} e^{-y/\theta_2}}{\frac{1}{2}e^{-y/2}} = \frac{2a_1}{\theta_1} e^{y(\frac{1}{2} - \frac{1}{\theta_1})} + \frac{2a_2}{\theta_2} e^{y(\frac{1}{2} - \frac{1}{\theta_2})}.$ Since  $\theta_1 < 2 < \theta_2$ , it follows that  $\frac{1}{\theta_2} < \frac{1}{2} < \frac{1}{\theta_1}$ . Therefore,  $\frac{1}{2} - \frac{1}{\theta_2} > 0$ so that  $\lim_{y \to \infty} \frac{f_Y(y)}{f_Z(y)} = \lim_{y \to \infty} \frac{2a_1}{\theta_1} e^{y(\frac{1}{2} - \frac{1}{\theta_1})} + \frac{2a_2}{\theta_2} e^{y(\frac{1}{2} - \frac{1}{\theta_2})} = 0 + \infty$ . True.

II. Suppose that  $f_Y(y) = \frac{1}{2}e^{-y} + \frac{1}{6}e^{-y/3}$ . Then  $S_Y(y) = \frac{1}{2}e^{-y} + \frac{1}{2}e^{-y/3}$ , so that Y is mixture of exponentials with means of 1 and 3, with equal mixing weights of  $\frac{1}{2}$ .  $e_Y(d) = \frac{\int_d^{\infty} S(y) \, dy}{g(x)} = \frac{\frac{1}{2}e^{-d} + \frac{3}{2}e^{-d/3}}{g(x)}$ .

$$e_{X_1}(d) = 1$$
,  $e_{X_3}(d) = 3 \rightarrow \frac{1}{2} \times e_{X_1}(d) + \frac{1}{2} \times e_{X_3}(d) = 2$ . False.

9. Since the two estimates are independent,

$$Var[\mu_C] = Var[w \times \mu_A + (1 - w) \times \mu_B] = w^2 \times Var[\mu_A] + (1 - w)^2 \times Var[\mu_B]$$
  
=  $w^2 \times 400^2 + (1 - w)^2 \times 200^2 = 200,000w^2 - 80,000w + 40,000.$ 

This is a quadratic expression in w with positive coefficient of  $w^2$ . The minimum can be found by differentiating with respect to w and setting equal to 0:

 $400,000w - 80,000 = 0 \rightarrow w = \frac{1}{5}$  is the value of w that minimizes  $Var[\mu_c]$ .

#### Answer B

10. If Y is a mixture of  $X_1$  and  $X_2$  with mixing weights a and 1-a, we can define the parameter  $\Theta = \{1,2\}$ , with  $P[\Theta = 1] = a$ ,  $P[\Theta = 2] = 1 - a$ .

Then  $E[Y] = E[E[Y|\Theta]] = E[Y|\Theta = 1] \times P[\Theta = 1] + E[Y|\Theta = 2] \times P[\Theta = 2]$ =  $E[X_1] \times a + E[X_2] \times (1 - a)$ 

(this is the usual way the mean of a finite mixture is formulated).

 $Var[Y] = E[Var[Y|\Theta]] + Var[E[Y|\Theta]], \text{ where}$   $E[Var[Y|\Theta]] = Var[X_1] \times a + Var[X_2] \times (1-a) \text{ and}$  $Var[E[Y|\Theta]] = (E[X_1] - E[X_2])^2 \times a \times (1-a).$ 

The last equality follows from the fact that if Z is a two-point random variable

$$Z = \begin{cases} u & \text{prob. } p \\ v & \text{prob. } 1 - p \end{cases}, \text{ then } Var[Z] = (u - v)^2 \times p \times (1 - p); E[Y|\Theta] \text{ is }$$

a two-point random variable  $E[Y|\Theta] = \begin{cases} E[X_1] & \text{prob. } a \\ E[X_2] & \text{prob. } 1-a \end{cases}$ 

Therefore  $Var[Y] = Var[X_1] \times a + Var[X_2] \times (1-a) + (E[X_1] - E[X_2])^2 \times a \times (1-a).$ 

B) If  $X_1, X_2$  are Poisson random variables then  $E[X_1] = Var[X_1]$  and  $E[X_2] = Var[X_2]$ , so that  $E[Y] = E[X_1] \times a + E[X_2] \times (1-a) = Var[X_1] \times a + Var[X_2] \times (1-a)$ .

It follows that  $Var[Y] = Var[X_1] \times a + Var[X_2] \times (1-a) + (E[X_1] - E[X_2])^2 \times a \times (1-a) > E[Y].$ 

C) For a negative binomial random variable with parameters r and  $\beta$ , the mean is  $r\beta$  and the variance is  $r\beta(1 + \beta)$ , so the variance is larger than the mean. If  $X_1$  and  $X_2$  have negative binomial distributions, the  $E[X_1] < Var[X_1]$  and  $E[X_2] < Var[X_2]$ .

Therefore,  $E[Y] = E[X_1] \times a + E[X_2] \times (1-a) < Var[X_1] \times a + Var[X_2] \times (1-a)$ , and  $Var[X_1] \times a + Var[X_2] \times (1-a)$   $< Var[X_1] \times a + Var[X_2] \times (1-a) + (E[X_1] - E[X_2])^2 \times a(1-a) = Var[Y]$ ; therefore E[Y] < Var[Y].

A) For a binomial random variable with parameters m and q, the mean is mq and the variance is mq(1-q), which is smaller than the mean.

Therefore  $E[Y] = E[X_1] \times a + E[X_2] \times (1-a) > Var[X_1] \times a + Var[X_2] \times (1-a)$ , and it is possible that when we add  $(E[X_1] - E[X_2])^2 \times a \times (1-a)$  to the right side, we get approximate equality.

So it is possible that E[Y] = Var[Y] for a mixture of binomials.

#### Answer A

11. The mean will be  $\frac{1}{2} \times (m_1 + m_2)$  and the variance will be  $\frac{1}{2} \times (2m_1^2 + 2m_2^2) - [\frac{1}{2} \times (m_1 + m_2)]^2$ . The coefficient of variation is the ratio of standard deviation to mean.

The square of the coefficient of variation is

$$\frac{\frac{1}{2} \times (2m_1^2 + 2m_2^2) - [\frac{1}{2} \times (m_1 + m_2)]^2}{[\frac{1}{2} \times (m_1 + m_2)]^2} = \frac{3m_1^2 - 2m_1m_2 + 3m_2^2}{(m_1 + m_2)^2}$$
$$= \frac{3m_1^2 + 6m_1m_2 + 3m_2^2 - 8m_1m_2}{(m_1 + m_2)^2}$$
$$= 3 - \frac{8m_1m_2}{(m_1 + m_2)^2}.$$

The maximum square of the coefficient of variation is 3, and it occurs at the minimum of  $\frac{8m_1m_2}{(m_1+m_2)^2}$ , which is 0 (if  $m_1$  or  $m_2$  is 0).

The least upper bound of the coefficient of variation is  $\sqrt{3}$ .

#### Answer C

#### **12.** Var[Z] = Var[E[Z | Y]] + E[Var[Z | Y]].

 $E[Z \mid Y] = 2 \times Y$  (a sum of Y independent normal random variables each with mean 2), and  $Var[Z \mid Y] = 2 \times Y$  (a sum of Y independent normal random variables each with a variance of 2).

Thus,  $Var[Z] = Var[2 \times Y] + E[2 \times Y] = 4 \times Var[Y] + 2 \times E[Y].$  Var[Y] = Var[E[Y | X]] + E[Var[Y | X]].Since E[Y] = 0.6 with prob. 0.4 (X = 0) and E[Y] = 1.3 with prob. 0.6 (X = 1), it follows that  $Var[E[Y | X]] = (0.6)^2 \times 0.4 + (1.3)^2 \times 0.6 - (1.02)^2 = 0.1176.$ 

Also, if X = 0, the variance of Y is  $1^2 \times 0.2 + 2^2 \times 0.2 - (0.6)^2 = 0.64$ , and if X = 1, the variance of Y is  $1^2 \times 0.3 + 2^2 \times 0.5 - (1.3)^2 = 0.61$ . Thus,  $E[Var[Y \mid X]] = 0.64 \times 0.4 + .61 \times .6 = 0.622$ , and so Var[Y] = 0.1176 + 0.622 = 0.7396.

Then, the variance of Z is  $4 \times 0.7396 + 2 \times 1.02 = 4.9984$ . Note that Var[Y] can also be found from  $E[Y^2] - (E[Y])^2$ . Again, E[Y] = 1.02, as above, and now,  $E[Y^2] = E[E[Y^2 | X]] = 1 \times 0.4 + 2.3 \times 0.6 = 1.78$ , so that Var[Y] = 0.7396.

#### Answer A

13. Given that a local train arrives first, you will get to work 28 minutes after that local train arrives, since you will take it. Your co-worker will wait for first express train. You will get to work before your co-worker if the next express train (after the local) arrives more than 12 minutes after the local. We expect 5 express trains per hour, so the time between express trains is exponentially distributed with a mean of  $\frac{1}{5}$  of an hour, or 12 minutes. Because of the lack of memory property of the exponential distribution, since we are given that the next train is local, the time until the next express train after that is exponential with a mean of 12 minutes. Therefore, the probability that after the local, the next express arrives in more than 12 minutes is P[T > 12], where T has an exponential distribution with a mean of 12. This probability is  $e^{-12/12} = e^{-1} = 0.368$  (37%).

#### Answer A



#### Ready for more practice? Check out GOAL!

GOAL offers additional questions, quizzes, and simulated exams with helpful solutions and tips. Included with GOAL are topic-by-topic instructional videos! Head to ActuarialUniversity.com and log into your account.





Do you have any fun ideas for future Study Breaks?

Share your inspiration with #actexstudybreak

Exercise is a proven way to boost your verbal memory, thinking, and learning. <sup>[1]</sup>



## **Pyramid with Arms Extended**

## Yoga - Sun Salutations

\*Hold each pose for three long breaths, breathing in through your nose and out through your mouth.

Note that we are mathematicians, not yogi: attempt poses at your own risk!



## Practice Exam 1

1. Clear Each of decision-makers X, Y, and Z has the opportunity to participate in a game with payoff uniformly distributed on (0, 10000). Assume that X, Y and Z value assets of amount  $w \ge 0$  according to the following utility functions:

Decision Maker	Utility Function
X	$u_X(w) = \sqrt{w}$
Y	$u_Y(w) = \frac{w}{100}$
Z	$u_Z(w) = \left(\frac{w}{100}\right)^2$

Which decision maker would not be willing to pay more than 5,000 to participate in the game?

(A) $X$ and $Y$ only	(B) $X$ and $Z$ only
(C) $Y$ and $Z$ only	(D) $X, Y$ and $Z$

- (E) The correct answer is not given by (A), (B), (C) or (D)
- 2. The XYZ Insurance Company sells property insurance policies with a deductible of \$5,000, policy limit of \$500,000, and a coinsurance factor of 80%. Let  $X_i$  be the individual loss amount of the *i*th claim and  $Y_i$  be the claim payment of the *i*th claim. Which of the following represents the relationship between  $X_i$  and  $Y_i$ ?

(A) $Y_i = \begin{cases} 0\\ 0.80 (X_i - 5,000)\\ 500,000 \end{cases}$	$X_i \le 5,000$ $5,000 < X_i \le 625,000$ $X_i > 625,000$
(B) $Y_i = \begin{cases} 0\\ 0.80 (X_i - 4,000)\\ 500,000 \end{cases}$	$X_i \le 5,000$ $4,000 < X_i \le 500,000$ $X_i > 500,000$
(C) $Y_i = \begin{cases} 0 \\ 0.80 (X_i - 5,000) \\ 500,000 \end{cases}$	$X_i \le 5,000$ $5,000 < X_i \le 630,000$ $X_i > 630,000$
(D) $Y_i = \begin{cases} 0\\ 0.80 (X_i - 6,250)\\ 500,000 \end{cases}$	$X_i \le 6,250$ $6,250 < X_i \le 631,500$ $X_i > 631,500$
(E) $Y_i = \begin{cases} 0\\ 0.80 (X_i - 5,000)\\ 500,000 \end{cases}$	$X_i \le 5,000$ $5,000 < X_i \le 505,000$ $X_i > 505,000$

- 3. A casino has a game that makes payouts at a Poisson rate of 5 per hour and the payout amounts can be any non-negative integer, 1, 2, 3, ... without limit. The probability that any given payout is equal to i > 0 is  $\frac{1}{2^i}$ . Payouts are independent. Calculate the probability that there are no payouts of 1,2, or 3 in a given 20 minute period.
  - (A) 0.08 (B) 0.13 (C) 0.18 (D) 0.23 (E) 0.28
- 4. Soom Buy Tire Store, a nationwide chain of retail tire stores, sells 2,000,000 tires per year of various sizes and models. Zoom Buy offers the following road hazard warranty: "If a tire sold by us is irreparably damaged in the first year after purchase, we'll replace it free, regardless of the cause."

The average annual cost of honoring this warranty is \$10,000,000, with a standard deviation of \$40,000. Individual claim counts follow a binomial distribution, and the average cost to replace a tire is \$100. All tires are equally likely to fail in the first year, and tire failures are independent. Calculate the standard deviation of the replacement cost per tire.

- (A) Less than 60
- (B) At least \$60, but less than \$65
- (C) At least 65, but less than 70
- (D) At least \$70, but less than \$75
- (E) At least \$75

$$egin{array}{ccc} x & f_X(x) \ 5 & 0.6 \ 10 & 0.4 \end{array}$$

Determine the expected cost of an aggregate stop-loss insurance with a deductible of 6.

- (A) Less than 15.0
- (B) At least 15.0 but less than 15.3  $\,$
- (C) At least 15.3 but less than 15.6
- (D) At least 15.6 but less than 15.9
- (E) At least 15.9

Use the following information for questions 6 and 7. You are the producer of a television quiz show that gives cash prizes. The number of prizes, N, and prize amounts, X are independent of one another and have the following distributions:

$$\begin{array}{ll} N: & P[N=1]=0.8, \quad P[N=2]=0.2 \\ X: & P[X=0]=0.2, \quad P[X=100]=0.7, \quad P[X=1000]=0.1 \end{array}$$

6. ♥ Your budget for prizes equals the expected prizes plus 1.25× standard deviation of prizes. Calculate your budget.

(A) 384 (B) 394 (C) 494 (D) 588 (E) 596

7. You buy stop-loss insurance for prizes with a deductible of 200. The cost of insurance includes a 175% relative security load (the relative security load is the percentage of expected payment that is added). Calculate the cost of the insurance.

(A) 204 (B) 227 (C) 245 (D) 273 (E) 357

- 8. •• An actuary determines that claim counts follow a negative binomial distribution with unknown  $\beta$  and r. It is also determined that individual claim amounts are independent and identically distributed with mean 700 and variance 1,300. Aggregate losses have a mean of 48,000 and variance 80 million. Calculate the values for  $\beta$  and r.
  - (A)  $\beta = 1.20, r = 57.19$ (B)  $\beta = 1.38, r = 49.75$ (C)  $\beta = 2.38, r = 28.83$ (D)  $\beta = 1,663.81, r = 0.04$
  - (E)  $\beta = 1,664.81, r = 0.04$
- 9. Clet  $X_1, X_2, X_3$  be independent Poisson random variables with means  $\theta, 2\theta$ , and  $3\theta$  respectively. What is the maximum likelihood estimator of  $\theta$  based on sample values  $x_1, x_2$ , and  $x_3$  from the distributions of  $X_1, X_2$  and  $X_3$ , respectively,
  - (A)  $\frac{\bar{x}}{2}$  (B)  $\bar{x}$ (C)  $\frac{x_1 + 2x_2 + 3x_3}{6}$  (D)  $\frac{3x_1 + 2x_2 + x_3}{6}$ (E)  $\frac{6x_1 + 3x_2 + 2x_3}{11}$

10. •• Loss random variable X has a uniform distribution on  $(0, \theta)$ . A sample is taken of n insurance payments from policies with a limit of 100. Eight of the sample values are limit payments of 100. The maximum likelihood estimate of  $\theta$  is  $\hat{\theta}$ . Another sample is taken, also of n insurance payments, but from policies with a limit of 150. Three of the sample values are limit payments of 150. The maximum likelihood estimate of  $\theta$  is  $\frac{4}{3}\hat{\theta}$ .

Determine n.

(A) 40  (B) 42  (C) 44  (D) 46	(E) 48
--------------------------------	--------

11. 🗣 For a group of policies, you are given:

- (i) Losses follow a uniform distribution on the interval  $(0, \theta)$ , where  $\theta > 25$ .
- (ii) A sample of 20 losses resulted in the following:

Interval	Number of Losses
$x \le 10$	$n_1$
$10 < x \le 25$	$n_2$
x > 25	$n_3$

The maximum likelihood estimate of  $\theta$  can be written in the form 25 + y. Determine y.

(A) 
$$\frac{25n_1}{n_2 + n_3}$$
  
(B)  $\frac{25n_2}{n_1 + n_3}$   
(C)  $\frac{25n_3}{n_1 + n_2}$   
(D)  $\frac{25n_1}{n_1 + n_2 + n_3}$   
(E)  $\frac{25n_2}{n_1 + n_2 + n_3}$ 

12. The number of claims follows a negative binomial distribution with parameters  $\beta$  and r, where  $\beta$  is unknown and r is known. You wish to estimate  $\beta$  based on n observations, where  $\bar{x}$  is the mean of these observations. Determine the maximum likelihood estimate of  $\beta$ .

(A) 
$$\frac{\bar{x}}{r^2}$$
 (B)  $\frac{\bar{x}}{r}$  (C)  $\bar{x}$  (D)  $r\bar{x}$  (E)  $r^2\bar{x}$ 

13. The following 6 observations are assumed to come from the continuous distribution with pdf  $f(x;\theta) = \frac{1}{2}x^2\theta^3 e^{-\theta x} : 1, 3, 4, 4, 5, 7.$ 

Find the mle of  $\theta$ .

(A) 
$$0.25$$
 (B)  $0.50$  (C)  $0.75$  (D)  $1.00$  (E)  $1.25$ 

14. An analysis of credibility premiums is being done for a particular compound Poisson claims distribution, where the criterion is that the total cost of claims is within 5% of the expected cost of claims with a probability of 90%. It is found that with n = 60 exposures (periods) and  $\bar{X} = 180.0$ , the credibility premium is 189.47. After 20 more exposures (for a total of 80) and revised  $\bar{X} = 185$ , the credibility premium is 190.88. After 20 more exposures (for a total of 100) the revised  $\bar{X}$  is 187.5. Assuming that the manual premium remains unchanged in all cases, and assuming that full credibility has not been reached in any of the cases, find the credibility premium for the 100 exposure case.

$$(A) 191.5 (B) 192.5 (C) 193.5 (D) 194.5 (E) 196.5 (E)$$

- 15. **V** For an insurance portfolio, you are given:
  - (i) For each individual insured, the number of claims follows a Poisson distribution.
  - (ii) The mean claim count varies by insured, and the distribution of mean claim counts follows gamma distribution.
  - (iii) For a random sample of 1000 insureds, the observed claim counts are as follows:

Number of Claims, $n$	0	1	2	3	4	5
Number of Insureds, $f_n$	512	307	123	41	11	6

$$\sum nf_n = 750, \qquad \sum n^2 f_n = 1494$$

- (iv) Claim sizes follow a Pareto distribution with mean 1500 and variance 6,750,000.
- (v) Claim sizes and claim counts are independent.
- (vi) The full credibility standard is to be within 5% of the expected aggregate loss 95% of the time.

Determine the minimum number of insureds needed for the aggregate loss to be fully credible.

- (A) Less than 8300 (B) At least 8300, but less than 8400
- (C) At least 8400, but less than 8500
- (D) At least 8500, but less than 8600

(E) At least 8600

Information on Questions 16 and 17 is as follows. You are given the following information on cumulative incurred losses through development years shown.

	Cumulative Incurred Losses				Paid-to-Date
Accident	Development Year			at Dec 31, AY4	
Year	0	1	2	3	
AY1	2325	3749	4577	4701	4701
AY2	2657	4438	5529		4500
AY3	2913	4995			3500
AY4	3163				2500

16. 😵 Using an average factor model, calculate the estimated total loss reserve as of Dec. 31, AY4.

(A) Less than 7500

- (B) At least 7500 but less than 7800
- (C) At least 7800 but less than 8100
- (E) At least 8400

(D) At least 8100 but less than 8400

17. 😪 As of Dec. 31, AY4, calculate

Estimated reserve for AY2 based on an average factor model) – (Estimated reserve for AY2 based on a mean factor model)

- (A) Less than -300
- (B) At least -300 but less than -100
- (C) At least -100 but less than 100
- (D) At least 100 but less than 300
- (E) At least 300
- 18. V For a one-period binomial model for the price of a stock with price 100 at time 0, you are given:
  - (i) The stock pays no dividends.
  - (ii) The stock price is either 110 or 95 at the end of the year.
  - (iii) The risk free force of interest is 5%.

Calculate the price at time 0 of a one-year call option with strike price 100.

- (A) Less than 6.00
- (B) At least 6.00 but less than 6.25
- (C) At least 6.25 but less than 6.50
- (D) At least 6.50 but less than 6.75
- (E) At least 6.75
- 19. Vising the following information, determine the incurred losses for the 2017 accident year as reported at Dec. 31, 2018.

Occurrence #1: Occurrence date Feb. 1/16, Report date Apr. 1/16

Loss History:

Date	Total Paid to Date	Unpaid Loss Reserve	Total Incurred
Apr. 1/16	1000	1000	2000
Dec. 31/16	1500	1000	2500
Dec. $31/17$	1500	1000	2500
Mar. 1/18	3000	0	3000

Date	Total Paid to Date	Unpaid Loss Reserve	Total Incurred
July 1/17	1000	2000	3000
Dec. 31/17	3000	1000	4000
Dec. 31/18	5000	0	5000

Occurrence #2: Occurrence date May 1/17, Report date July 1/17 Loss History:

Occurrence #3: Occurrence date Nov. 1/17, Report date Feb. 1/18 Loss History

Date	Total Paid to Date	Unpaid Loss Reserve	Total Incurred
Mar. 1/18	0	8000	8000
Dec. 31/18	5000	5000	10,000
(A) 0		(B) 3,00	0
(C) 5,000	)	(D) 8,00	0
(E) 15,00	00		

20.  $\checkmark$  You are given the following calendar year earned premium.

Year	CY2	CY3	CY4		
Earned Premium	4200	4700	5000		
You are also given the	following r	ate chang	es		
Date	April	1, CY1	Septe	mber 1, CY2	July 1, CY3
Average Rate Change	+1	12~%		+6 %	+10 %
Determine the approximate earned premium at current (end of CY4) rates for CY3.					
(A) Less than 5000 (B) At least 5000 but less than 510				but less than 5100	
(C) At least 5100 but less than 5200 (D) At least 5200 but less than 53 $($			but less than $5300$		
(E) At least $5300$					

#### Solutions to Practice Exam 1

1.	А	
2.	С	
3.	D	
4.	Е	
5.	С	
6.	Е	
7.	D	
8.	В	
9.	А	
10.	Е	
11.	С	
12.	В	
13.	С	
14.	А	
15.	Е	
16.	D	
17.	С	
18.	С	
19.	Е	
20.	С	

1. A decision maker with utility function u(w) will pay amount C to play a game (random variable) W based on the following relationship: u(C) = E[u(W)].

For individual X:  $E[u_X(W)] = \int_0^{10,000} \frac{\sqrt{w}}{10,000} dw = \frac{200}{3} = u_X(C_X) = \sqrt{C_X}.$ Solving for  $C_X$  results in  $C_X = \left(\frac{200}{3}\right)^2 = 4,444.44$ 

For individual Y:  $E[u_Y(W)] = \int_0^{10,000} \frac{w}{100} \times \frac{1}{10,000} dw = 50 = u_Y(C_Y) = \frac{C_Y}{100}$ . Solving for  $C_Y$  results in  $C_Y = 5,000$ 

For individual Z:  $E[u_Z(W)] = \int_0^{10,000} \left(\frac{w}{100}\right)^2 \times \frac{1}{10,000} dw = \frac{10,000}{3} = u_Z(C_Z) = \left(\frac{C_Z}{100}\right)^2$ . Solving for  $C_Z$  results in  $C_Z = 100 \times \sqrt{\frac{10,000}{3}} = 5,773.50$ 

Only Z will pay more than 5,000 for the gamble. In general the concavity of the utility function determines the risk profile of the individual. X has a concave utility function  $(u''_X(w) < 0)$ and will be risk-averse and will not be willing to pay more that the expected value (fair cost) of the gamble of 5,000. Y has a linear utility function and will be risk-neutral. Y will be willing to pay at most the expected payoff of the gamble of 5.000 but not more. Z has a convex utility function  $(u''_X(w) > 0)$  and will be risk-preferring and will pay more the 5,000 for the gamble.

#### Answer A

2. With coinsurance factor  $\alpha$ , deductible d, policy limit  $\alpha(u-d)$ , the amount paid per loss is

(we are assuming in inflation rate of 
$$r = 0$$
) $Y = \begin{cases} 0 & X \le d \\ \alpha(X - d) & d < X \le u. \\ \alpha(u - d) & X > u \end{cases}$ 

In this problem, the coinsurance factor is  $\alpha = .8$ , the deductible is d = 5,000, and the policy limit is .8(u - 5,000) = 500,000, so that the maximum covered loss is u = 630,000.

The amount paid per loss becomes 
$$Y = \begin{cases} 0 & X \le 5,000 \\ 0.80(X - 5,000) & 5,000 < X \le 630,000 \\ 500,000 & X > 630,000 \end{cases}$$
 Answer C

3. When a payout occurs, it is 1, 2 or 3 with probability  $\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} = \frac{7}{8}$ . The number of payouts that are 1,2 or 3 follows a Poisson process with an hourly rate of  $5 \times \frac{7}{8} = \frac{35}{8}$ . The expected number of payouts that are 1, 2 or 3 in 20 minutes, say N, has a Poisson distribution with mean  $\frac{35}{8} \times \frac{20}{60} = \frac{35}{24}$ . The probability that there are no payouts of 1, 2, or 3 in a given 20 minute period is the probability that N = 0, which is  $e^{-35/24} = .233$ .

#### Answer D